


Easy Steps for Optimal Yeast Rehydration

Proper yeast rehydration is one of the most important steps to help ensure a strong and healthy fermentation. Normal inoculation for wine active dried yeast is 2 lb/1000 gal (25 g/hL). When added properly, a 2 lb/1000 gal (25 g/hL) addition of wine active dried yeast results in an initial cell concentration of 3-4 million viable cells per milliliter of must/juice. Under favorable conditions, the initial cell population may increase up to 100-150 million viable cells per milliliter of must/juice before growth stops and alcoholic fermentation begins. This biomass increase is critical for healthy fermentations. When harvesting grapes at high maturity levels, increased inoculation rates are recommended. Careful rehydration, atemperation and inoculation are all important to help prevent sluggish or stuck fermentations.



1

Suspend 2.5 lb/1000 gal (30g/hL) of an appropriate yeast rehydration in 20 times its weight of clean, chlorine free 43°C(110°F) water. If the water temperature is not high enough, the yeast rehydration nutrient may not go entirely into solution. We recommend Lalvin GoFerm and Laffort Dynastart as appropriate yeast rehydration nutrients. **IMPORTANT:** If not using a yeast rehydration nutrient, water temperature should begin at 40°C(104°F) to avoid harming the yeast.

2

Once the temperature of the yeast rehydration nutrient solution has dropped to 40°C(104°F), add 2 lb/1000 gal (25 g/hL) of active dried yeast. Stir gently to break up any clumps. Let suspension stand for 15-30 minutes, then stir gently again. Allowing rehydrated yeast to stand for more than 30 minutes may start a decline in the live population. **Note:** Foaming is not an indicator of yeast viability.

3

Slowly (over a period of 5 minutes) combine an equal amount of the must/juice to be fermented with the yeast suspension. This will help the yeast adjust to the cool temperature of the must/juice and will help avoid cold shock caused by a rapid temperature drop exceeding 10°C(18°F). This atemperation step may need repeating in very low temperature must. For every 10°C(18°F) temperature difference between the must/juice and the yeast slurry, an atemperation step must be performed. For example, for a must/juice temperature of 20°C(68°F) and yeast slurry temperature of 40°C(104°F), two atemperation steps are required.

4

Add the yeast slurry to the bottom of the fermentation vessel just as you begin filling the vessel with must/juice. This is especially important for large tanks with long filling times or when inoculating with strains that are sensitive to the competitive factor. This will allow the yeast a head start over indigenous organisms.

Cuatro pasos fáciles para la rehidratación de la levadura y la inoculación del mosto

Yeast Rehydration Nutrient



1

Suspender 2.5lb/1000gal (30 g/hl) de un nutriente de rehidratación en 20 veces su peso en agua limpia 110°F (43°C). **IMPORTANTE:** Si no se utiliza un nutriente de rehidratación, la temperatura del agua debe ser 104°F (40°C) para evitar dañar la levadura.




2

Una vez que la temperatura de la solución con un nutriente de rehidratación ha descendido hasta 104°F (40°C), añadir 2lb/1000gal (25 g/hl) de levaduras seleccionadas. Agitar suavemente para eliminar cualquier grumo. Dejar reposar 15-30 minutos y agitar de nuevo lentamente.



3

Mezclar lentamente durante 5 minutos la levadura en suspensión con una cantidad igual de mosto (emplear el mosto que se va a inocular). Esto ayudará a la levadura a adaptarse a temperaturas más bajas y se evitará el choque térmico provocado por una bajada rápida de la temperatura (evitar una diferencia de más de 18°F o 10°C). Puede ser necesario repetir esta aclimatación en mostos de temperatura muy baja.



4

Adicionar la levadura en el fondo del depósito de fermentación tan pronto como se empiece a llenar con mosto o uvas.


Quatre étapes faciles pour réhydrater les levures et inoculer le moût

Yeast Rehydration Nutrient



1

Ajouter 30 g de aliment de réhydratation par hl de moût à fermenter, dans 20 fois le poids de aliment de réhydratation en eau propre (à 43°C).
IMPORTANT: Si vous n'utilisez pas aliment de réhydratation, l'eau doit être à 40°C pour éviter une baisse de population de levures.




2

Une fois le température de la suspension de aliment de réhydratation descendue à 40°C, ajouter 25 g/hl de levures sélectionnées à la suspension de aliment de réhydratation. Remuer doucement pour éviter les grumeaux. Laisser reposer sans remuer pendant 15-30 minutes, ensuite bien agiter la solution pour obtenir une bonne suspension.



3

Ajouter lentement (5 minutes) un volume égal du moût à ensemer. Pour les moûts à très basse température, cette opération devra alors être répétée afin d'éviter le choc thermique qui se produit quand le différentiel de température entre le moût et la suspension de levures est supérieur à 10°C.



4

Ajouter la suspension de levures au fond de la cuve pendant que vous commencez à ajouter le moût.